

# Temporal Variability of Urinary Phthalate Metabolite Levels in Men of Reproductive Age

Russ Hauser, John D. Meeker, Sohee Park, Manori J. Silva,  
and Antonia M. Calafat

doi:10.1289/ehp.7212 (available at <http://dx.doi.org/>)

Online 16 August 2004



“Temporal Variability of Urinary Phthalate Metabolite Levels in Men of Reproductive Age”

Russ Hauser<sup>a,b</sup>, John D. Meeker<sup>a</sup>, Sohee Park<sup>c</sup>, Manori J. Silva<sup>d</sup>, Antonia M. Calafat<sup>d</sup>.

<sup>a</sup> Department of Environmental Health, Harvard School of Public Health, Boston, MA.

<sup>b</sup> Vincent Memorial Obstetrics & Gynecology Service, Andrology Laboratory and In Vitro Fertilization Unit, Massachusetts General Hospital, Boston, MA 02114.

<sup>c</sup> Department of Biostatistics, Harvard School of Public Health, Boston, MA.

<sup>d</sup> National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA.

Correspondence should be directed to:

Russ Hauser MD, ScD, MPH

Harvard School of Public Health

Department of Environmental Health

Occupational Health Program

Building 1- room 1405

665 Huntington Avenue

Boston, MA 02115

(office) 617-432-3326

(fax) 617-432-0219

(e-mail) [rhauser@hohp.harvard.edu](mailto:rhauser@hohp.harvard.edu)

Running title: Variability of urinary phthalates.

Key words: Biomarkers, human, phthalates, reliability, urine.

#### Acknowledgments:

The authors thank Dr. J. Reidy, Ms. A. Herbert, Ms. E. Samandar, and Mr. J. Preau from the CDC and Dr. S. Duty, Ms. J. Frelich, Ms. L. Godfrey-Bailey, Ms. A. Trisini and Mr. R. Dadd from HSPH.

Supported by grant numbers ES09718 and ES00002 from the National Institute of Environmental Health Sciences (NIEHS), NIH. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NIEHS.

#### Abbreviations:

BBzP:	Butyl benzyl phthalate.
CDC:	Centers for Disease Control and Prevention.
DBP:	Di-n-butyl phthalate.
DEHP:	Di-2-ethylhexyl phthalate.
DEP:	Di-ethyl phthalate.
HSPH:	Harvard School of Public Health.
ICCs:	Intra-class correlation coefficients.
LOD:	Limit of detection.
MBP:	Mono-butyl phthalate.
MBzP:	Mono-benzyl phthalate.
MCHP:	Monocyclohexyl phthalate.
MEHP:	Mono-2-ethylhexyl phthalate.
MEP:	Mono-ethyl phthalate.

MGH:	Massachusetts General Hospital.
MINP:	Mono-3-methyl-5-dimethylhexyl phthalate.
MMP:	Mono-methyl phthalate.
MOP:	Mono-n-octyl phthalate.
PVC:	Polyvinyl chloride.
SAS:	Statistical Analysis Software.
µg/L:	Micrograms per liter.

## OUTLINE

ABSTRACT

INTRODUCTION

METHODS

STATISTICAL ANALYSES

RESULTS

DISCUSSION

REFERENCES

TABLES 1-4

FIGURES 1-5

## Abstract

Phthalates are a family of multifunctional chemicals widely used in personal care and other consumer products. The ubiquitous use of phthalates results in human exposure through multiple sources and routes, including dietary ingestion, dermal absorption, inhalation, as well as through parenteral exposure from medical devices containing phthalates. We explored the temporal variability over three-months in urinary phthalate metabolite levels among eleven men who collected up to nine urine samples each during this time period. Eight phthalate metabolites were measured by solid phase extraction-high performance liquid chromatography-tandem mass spectrometry. Statistical analyses were performed to determine the between- and within-subject variance apportionment, and the sensitivity and specificity of a single urine sample to classify a subject's long-term average exposure. Five of the eight phthalates were frequently detected. Mono-ethyl phthalate (MEP) was detected in 100 percent of samples; mono-butyl phthalate, mono-benzyl phthalate, mono-2-ethylhexyl phthalate (MEHP) and mono-methyl phthalate were detected in over 90 percent of samples. Although we found both substantial day-to-day and month-to-month variability in each individual's urinary phthalate metabolite levels, a single urine sample was moderately predictive of each subject's exposure over three months. The sensitivities ranged from 0.56 to 0.74. Both the degree of between- and within-subject variance and the predictive ability of a single urine sample differed among phthalate metabolites. A single urine sample was moderately predictive of 3-month exposure, although the predictive ability varied across phthalate monoesters. In particular, a single urine sample was most predictive for MEP and least predictive for MEHP. These results suggest that the most efficient exposure assessment strategy for a particular study may depend on the phthalates of interest.